

The large CP phase in $B_s - \bar{B}_s$ mixing from primary scalar unparticles

J. K. Parry

*Kavli Institute for Theoretical Physics China,
Institute of Theoretical Physics,
Chinese Academy of Science, Beijing 100190, China
and
Center for High Energy Physics,
Tsinghua University, Beijing 100084, China*

In this letter we consider the case of primary scalar unparticle contributions to $B_{d,s}$ mixing. With particular emphasis on the impact of the recent hint of new physics in the measurement of the B_s mixing phase, ϕ_s , we determine the allowed parameter space and impose bounds on the unparticle couplings.

PACS: 13.20.He 13.25.Hw 12.60.-i

I. INTRODUCTION

It has recently been suggested [1, 2] that there may exist a non-trivial scale invariant sector at high energies, known as unparticle stuff. These new fields with an infrared fixed point are called Banks-Zaks fields [3], interacting with Standard model fields via heavy particle exchange,

$$\frac{1}{M_{\mathcal{U}}^k} \mathcal{O}_{\text{SM}} \mathcal{O}_{BZ}. \quad (1)$$

Here $\mathcal{O}_{\text{SM}}$ is a standard model(SM) operator of mass dimension d_{SM} , $\mathcal{O}_{BZ}$ is a Banks-Zaks($\mathcal{BZ}$) operator of mass dimension $d_{\mathcal{BZ}}$, with $k = d_{\text{SM}} + d_{\mathcal{BZ}} - 4$, and $M_{\mathcal{U}}$ is the mass of the heavy particles mediating the interaction. At a scale denoted by $\Lambda_{\mathcal{U}}$ the $\mathcal{BZ}$ operators match onto unparticle operators with a new set of interactions,

$$C_{\mathcal{U}} \frac{\Lambda_{\mathcal{U}}^{d_{\mathcal{BZ}} - d_{\mathcal{U}}}}{M_{\mathcal{U}}^k} \mathcal{O}_{\text{SM}} \mathcal{O}_{\mathcal{U}} \quad (2)$$

where $\mathcal{O}_{\mathcal{U}}$ is an unparticle operator with scaling dimension $d_{\mathcal{U}}$ and $C_{\mathcal{U}}$ is the coefficient of the low energy theory. Unparticle stuff of scaling dimension $d_{\mathcal{U}}$ looks like a non-integral number $d_{\mathcal{U}}$ of invisible massless particles. It was recently suggested in [27] that conformal invariance implies constraints on the scaling dimension, $1 \leq d_{\mathcal{U}} \leq 2$, and $3 \leq d_{\mathcal{U}} \leq 4$, for scalar and vector unparticles respectively. This also implies the constraint $2 \leq d_{\mathcal{U}} \leq 3$ for the non-primary operator $\partial^\mu \mathcal{O}_{\mathcal{U}}$. We shall see later that assuming such constraints leads to the contributions from vector unparticles and from the non-primary operator $\partial^\mu \mathcal{O}_{\mathcal{U}}$ being negligible, except in a small region of parameter space where the scaling dimension approaches integer values. As a result, in this work we shall follow the above suggestion and focus on the case of the primary scalar unparticle ($\mathcal{O}_{\mathcal{U}}$) with couplings to the SM quarks as follows,

$$\mathcal{L}_{S'} = \frac{c_L^{S', q'q}}{\Lambda_{\mathcal{U}}^{d_{\mathcal{U}}-1}} \bar{q}'(1 - \gamma_5)q \mathcal{O}_{\mathcal{U}} + \frac{c_R^{S', q'q}}{\Lambda_{\mathcal{U}}^{d_{\mathcal{U}}-1}} \bar{q}'(1 + \gamma_5)q \mathcal{O}_{\mathcal{U}} \quad (3)$$

Here we assume that the left-handed and right-handed flavour-dependent dimensionless couplings $c_L^{S', bq}$, $c_R^{S', bq}$, are independent parameters. The above notation, $\mathcal{L}_{S', bq}$, has been used to distinguish the primary scalar operator studied here from $\mathcal{L}_S$ for the non-primary scalar case $\partial^\mu \mathcal{O}_{\mathcal{U}}$ studied previously [13]. We analyze a number of scenarios, in each case determining the allowed parameter space and placing bounds on the unparticle couplings.

The propagators for scalar unparticle are as follows [2, 4],

$$\int d^4x e^{iP \cdot x} \langle 0 | T \mathcal{O}_{\mathcal{U}}(x) \mathcal{O}_{\mathcal{U}}(0) | 0 \rangle = i \frac{A_{d_{\mathcal{U}}}}{2 \sin d_{\mathcal{U}} \pi} \frac{1}{(P^2 + i\epsilon)^{2-d_{\mathcal{U}}}} e^{-i\phi_{\mathcal{U}}} \quad (4)$$

$$(5)$$

where

$$A_{d_{\mathcal{U}}} = \frac{16\pi^{5/2}}{(2\pi)^{2d_{\mathcal{U}}}} \frac{\Gamma(d_{\mathcal{U}} + 1/2)}{\Gamma(d_{\mathcal{U}} - 1)\Gamma(2d_{\mathcal{U}})}, \quad \phi_{\mathcal{U}} = (d_{\mathcal{U}} - 2)\pi \quad (6)$$

The effects of unparticles, both scalar and vector, on meson-anti-meson mixing has been studied in the literature [5, 6, 7, 8, 9, 10, 11, 12, 13, 14]. In this work we shall study the constraints on primary scalar unparticles coming from the measurements of $B_{s,d}$ meson mass differences $\Delta M_{s,d}$ and also their CP violating phases $\phi_{s,d}$. In the B_d system these quantities have been well measured for some time and show no sizeable deviations from the SM expectations. In the B_s system recent measurements have also found small discrepancies between the SM expectation for ΔM_s [15], but now the CP violating phase ϕ_s , measured by the DØ[16] and CDF[17] collaborations reveals a deviation of 3σ [18, 19] [29]. This is the first evidence for new physics in $b \leftrightarrow s$ transitions. Studying primary scalar unparticles, we derive the constraints imposed by these latest measurements on the coupling between SM fields and unparticles, with particular interest on the impact of ϕ_s .

II. MESON-ANTIMESON MIXING FROM UNPARTICLES

Using the interactions listed in eq. (3) to evaluate the s- and t-channel contributions to meson mixing we obtain the effective Hamiltonian,

$$\mathcal{H}_{\text{eff}}^{S',q'q} = \frac{A_{d\mathcal{U}}}{2 \sin d\mathcal{U} \pi} \frac{e^{-i\phi_{\mathcal{U}}}}{\Lambda_{\mathcal{U}}^{2d_{\mathcal{U}}-2}} \left(\frac{1}{t^2-d_{\mathcal{U}}} + \frac{1}{s^2-d_{\mathcal{U}}} \right) \left[Q_2 \left(c_L^{S',q'q} \right)^2 + \tilde{Q}_2 \left(c_R^{S',q'q} \right)^2 + 2 Q_4 \left(c_L^{S',q'q} c_R^{S',q'q} \right) \right] \quad (7)$$

Here we have defined the quark operators $Q_1 - Q_5$ and their hadronic matrix elements as in [13]. Writing the $\Delta F = 2$ effective Hamiltonian in terms of these operators we have,

$$\mathcal{H}_{\text{eff}}^{q'q} = \sum_{i=1}^5 C_i^{S'} Q_i + \sum_{j=1}^3 \tilde{C}_j^{S'} \tilde{Q}_j \quad (8)$$

Here the operators $\tilde{Q}_{1,2,3}$ are obtained from $Q_{1,2,3}$ by the exchange $L \leftrightarrow R$. From eq. (7), it is straightforward to calculate the Wilson coefficients for primary scalar unparticles which are as follows,

$$C_2^{S'} = \frac{A_{d\mathcal{U}}}{\sin d\mathcal{U} \pi} \frac{e^{-i\phi_{\mathcal{U}}}}{M_M^2} \left(\frac{M_M^2}{\Lambda_{\mathcal{U}}^2} \right)^{d_{\mathcal{U}}-1} \left(c_L^{S',q'q} \right)^2 \quad (9)$$

$$C_4^{S'} = 2 \frac{A_{d\mathcal{U}}}{\sin d\mathcal{U} \pi} \frac{e^{-i\phi_{\mathcal{U}}}}{M_M^2} \left(\frac{M_M^2}{\Lambda_{\mathcal{U}}^2} \right)^{d_{\mathcal{U}}-1} \left(c_L^{S',q'q} c_R^{S',q'q} \right) \quad (10)$$

$$\tilde{C}_2^{S'} = \frac{A_{d\mathcal{U}}}{\sin d\mathcal{U} \pi} \frac{e^{-i\phi_{\mathcal{U}}}}{M_M^2} \left(\frac{M_M^2}{\Lambda_{\mathcal{U}}^2} \right)^{d_{\mathcal{U}}-1} \left(c_R^{S',q'q} \right)^2 \quad (11)$$

$$C_1^{S'} = C_3^{S'} = C_5^{S'} = \tilde{C}_1^{S'} = \tilde{C}_3^{S'} = 0 \quad (12)$$

where we have approximated $t = s \sim M_M^2$. This set of Wilson coefficients is rather similar to that for the non-primary operator $\partial^\mu \mathcal{O}_{\mathcal{U}}$, with the main differences being the sign of C_2 and the power to which the unparticle scale $\Lambda_{\mathcal{U}}$ is raised. These Wilson coefficients will mix with each other as a result of renormalisation group(RG) running down to the scale of M_M . For the B system, with a scale of new physics $\Lambda_{\mathcal{U}} = 1$ TeV, these Wilson coefficients at the scale $\mu_b = m_b$ are approximated as in [13]. The $\Delta F = 2$ transitions are defined as,

$$\langle \overline{M}^0 | \mathcal{H}_{\text{eff}}^{\Delta F=2} | M^0 \rangle = M_{12} \quad (13)$$

with the meson mass eigenstate difference defined as, $\Delta M \equiv M_H - M_L = 2|M_{12}|$. We can define in a model independent way the contribution to meson mixings in the presence of New Physics(NP) as,

$$M_{12} = M_{12}^{\text{SM}}(1 + R) \quad (14)$$

where M_{12}^{SM} denotes the SM contribution and $R \equiv r e^{i\sigma} = M_{12}^{\text{NP}}/M_{12}^{\text{SM}}$ parameterizes the NP contribution. The associated CP phase may then be defined as,

$$\phi \equiv \arg(M_{12}) = \phi^{\text{SM}} + \phi^{\text{NP}} \quad (15)$$

where $\phi^{\text{SM}} = \arg(M_{12}^{\text{SM}})$ and $\phi^{\text{NP}} = \arg(1 + r e^{i\sigma})$.

A. $B_{s,d}$ mixing and unparticle physics

In this work we shall focus on the constraints imposed on unparticle physics couplings from $B_{s,d}$ mixing. Therefore we set $q' = b, q = s, d$ and $M^0 = B_s^0, B_d^0$. In the B system, the Standard Model contribution to M_{12}^q is given by,

$$M_{12}^{q,\text{SM}} = \frac{G_F^2 M_W^2}{12\pi^2} M_{B_q} \hat{\eta}^B f_{B_q}^2 \hat{B}_{B_q} (V_{tq}^* V_{tb})^2 S_0(x_t) \quad (16)$$

where G_F is Fermi's constant, M_W the mass of the W boson, $\hat{\eta}^B = 0.551$ [22] is a short-distance QCD correction identical for both the B_s and B_d systems. The bag parameter $\hat{B}_{B_q}$ and decay constant f_{B_q} are non-perturbative quantities and contain the majority of the theoretical uncertainty. V_{tq} and V_{tb} are elements of the Cabibbo-Kobayashi-Maskawa (CKM) matrix [23, 24], and $S_0(x_t \equiv \bar{m}_t^2/M_W^2) = 2.34 \pm 0.03$, with $\bar{m}_t(m_t) = 164.5 \pm 1.1$ GeV [25], is one of the Inami-Lim functions [26].

We can now constrain both the magnitude and phase of the NP contribution, r_q and σ_q , through the comparison of the experimental measurements with SM expectations. From the definition of eq. (14) we have the constraint,

$$\rho_q \equiv \frac{\Delta M_q}{\Delta M_q^{\text{SM}}} = \sqrt{1 + 2r_q \cos \sigma_q + r_q^2} \quad (17)$$

The values for ρ_q given by the UTfit analysis [18, 20] at the 95% C.L. are,

$$\rho_d = [0.53, 2.05], \quad \rho_s = [0.62, 1.93] \quad (18)$$

These constraints on ρ_q encode the CP conserving measurements of $\Delta M_{d,s}$. The phase associated with NP can also be written in terms of r_q and σ_q ,

$$\sin \phi_q^{\text{NP}} = \frac{r_q \sin \sigma_q}{\sqrt{1 + 2r_q \cos \sigma_q + r_q^2}} \quad (19)$$

Here [18, 20] gives the 95% C.L. constraints,

$$\phi_d^{\text{NP}} = [-16.6, 3.2]^\circ \quad (20)$$

$$\phi_s^{\text{NP}} = [-156.90, -106.40]^\circ \cup [-60.9, -18.58]^\circ \quad (21)$$

these constraint represent those of the CP phase measurements of $\phi_{d,s}$. As in [13], in order to consistently apply the above constraints all input parameters are chosen to match those used in the analysis of the UTfit group [18, 20].

III. NUMERICAL ANALYSIS

First let us re-examine the contributions to $B_{d,s}$ mixing from primary vector unparticles $\mathcal{O}_U^\mu$ and the non-primary $\partial^\mu \mathcal{O}_U$. The Wilson coefficients for these cases can be found for example in [13]. Following the arguments of [27] we have the bounds $2 \leq d_U \leq 3$ for $\partial^\mu \mathcal{O}_U$ and $3 \leq d_U \leq 4$ for $\mathcal{O}_U^\mu$. These bounds produce a very large suppression of the contributions to $B_{d,s}$ mixing from these unparticle operators, unless we are sufficiently close to the pole in the unparticle propagator. These poles occur as d_U approaches integer values, due the $1/\sin d_U \pi$ nature of the propagator. Therefore the only significant contributions may occur for values of the scaling dimension, $d_U = A + \epsilon$ and $d_U = B - \delta$, where $A = 2(3)$ and $B = 3(4)$ for $\partial^\mu \mathcal{O}_U(\mathcal{O}_U^\mu)$, with $\epsilon, \delta \ll 1$. Fig. 1 and 2 show the size of ρ_s which results from varying ϵ and δ for couplings between SM operator and unparticle operator in the range $(-1, 1)$. From these plots we can see that for the unparticle operator $\partial^\mu \mathcal{O}_U$, we require $\epsilon \lesssim 0.1$ or $\delta \lesssim 10^{-7}$ to get significant contributions to B_s mixing. For the vector unparticle operator $\mathcal{O}_U^\mu$ the requirement is $\epsilon \lesssim 0.0005$ or $\delta \lesssim 10^{-10}$. The range of values of d_U for which the present allowed region for ρ_s may provide a constraint on the couplings is even smaller than those stated above. For example for the non-primary operator $\partial^\mu \mathcal{O}_U$ we require $\epsilon \lesssim 0.025$ before the ρ_s constraints begin to have any effect on the size of the couplings. The requirements for non-negligible contributions to B_d mixing will be less strict. It seems that, barring a small region of parameter space, the contributions from primary vector and non-primary scalar unparticles are negligible. As a result we here consider only the primary scalar unparticle contribution to $B_{s,d}$ mixing.

For our analysis of the primary scalar contribution to $B_{s,d}$ mixing we shall take the unparticle scale $\Lambda_U = 1$ TeV and generally fix $d_U = \frac{3}{2}$. In the following we shall consider possible coupling patterns as follows,




FIG. 1: Size of ρ_s against ϵ (left) and δ (right) for unparticle operator $\partial^\mu \mathcal{O}_U$ with real couplings $c_L^{S',bs}, c_R^{S',bs} \in (-1, 1)$ and unparticle scale $\Lambda_U = 1$ TeV.




FIG. 2: Size of ρ_s against ϵ (left) and δ (right) for unparticle operator $\mathcal{O}_U^\mu$ with real couplings $c_L^{V,bs}, c_R^{V,bs} \in (-1, 1)$ and unparticle scale $\Lambda_U = 1$ TeV.

- One real coupling; $c_L^{S',bq} \neq 0$ and $c_R^{S',bq} = 0$, with $c_L^{S',bq} \in \mathbb{R}$
- Two real couplings; $c_L^{S',bq} \neq 0$ and $c_R^{S',bq} \neq 0$, with $\{c_L^{S',bq}, c_R^{S',bq}\} \in \mathbb{R}$
- One complex coupling; $c_L^{S',bq} \neq 0$ and $c_R^{S',bq} = 0$, with $c_L^{S',bq} \in \mathbb{C}$
- Two complex couplings; $c_L^{S',bq} \neq 0$ and $c_R^{S',bq} \neq 0$, with $\{c_L^{S',bq}, c_R^{S',bq}\} \in \mathbb{C}$

In the first case, a single real coupling ($c_L^{S',bq} \neq 0, c_R^{S',bq} = 0$) between primary scalar unparticle operator and our SM quark operator, we allow the scaling dimension to vary in the range indicated by the bounds discussed earlier. Fig. 3 and 4 show the experimentally allowed parameter space in this case in the plane of the scaling dimension, d_U , and the coupling $c_L^{S',bq}$. This plot shows a very similar behaviour to that found for non-primary scalar unparticles with real couplings $c_L^{S,bq} = c_R^{S,bq}$ [13], with the allowed parameter space generally increasing for increasing d_U . This similarity between these two seemingly different cases is simply because they both give negative M_{12}^U due to the sign of the Wilson coefficient C_2 .

Fig. 3 shows the allowed $d_U - c_L^{S',bq}$ parameter space, black points obey the constraint from ΔM_d while grey points obey constraints from both ΔM_d and ϕ_d . From the left panel of fig. 3, with $d_U = \frac{3}{2}$, we can extract bounds,

$$|c_L^{S',bd}| \leq 0.0002 \quad (\Delta M_d \text{ only}) \quad (22)$$

$$|c_L^{S',bd}| \leq 0.00005 \quad (\Delta M_d \& \phi_d) \quad (23)$$




FIG. 3: Constraints on the $d_{\mathcal{U}}$ versus $c_L^{S',bq}$ parameter space from B_d mixing(left) and B_s mixing(right) for the case of a single real coupling $c_L^{S',bq} \neq 0$ and $c_R^{S',bq} = 0$. Black points indicate the $\Delta M_{d,s}$ allowed regions, while grey points indicate the regions are in agreement with both ΔM_q and the CP phase ϕ_q .

From the right panel of fig. 3, with $d_{\mathcal{U}} = \frac{3}{2}$, we can extract the bounds,

$$|c_L^{S',bs}| \leq 0.001 \quad (\Delta M_s \text{ only}) \quad (24)$$

$$|c_L^{S',bs}| \text{ excluded} \quad (\Delta M_s \& \phi_s) \quad (25)$$

where we see that the combined constraints of ΔM_s & ϕ_s exclude this possibility at the 3σ level, for all values of $d_{\mathcal{U}}$.




FIG. 4: Plot of the allowed $c_L^{S',bq}$ - $c_R^{S',bq}$ parameter space for B_d mixing(left) and B_s mixing(right) in the case of two real couplings and scaling dimension fixed as $d_{\mathcal{U}} = 3/2$. Black points show regions which agree with the measurement of $\Delta M_{d,s}$ while grey points show additional agreement with the measurement of the CP phases $\phi_{d,s}$.

For the second case of two real couplings, $c_L^{S',bq}$ and $c_R^{S',bq}$, we take two sub-cases, $c_L^{S',bq} \neq c_R^{S',bq}$ and $c_L^{S',bq} = c_R^{S',bq}$, shown in fig. 4 and 5 respectively. For the first sub-case, $c_L^{S',bq} \neq c_R^{S',bq}$, shown in fig. 4, there are generally no bounds that can be set on the couplings with the allowed parameter space stretching along two lines, similar to the case of two real non-equal couplings to unparticle operator $\partial^\mu \mathcal{O}_{\mathcal{U}}$. Looking at the grey area of fig. 4 we can see that including the ϕ_s and ϕ_d constraints further restrict the allowed parameter space. For the second sub-case, $c_L^{S',bq} = c_R^{S',bq}$, shown in fig. 5, the allowed parameter space is rather similar to the case of unparticle operators $\mathcal{O}_{\mathcal{U}}^\mu$ and $\partial^\mu \mathcal{O}_{\mathcal{U}}$ with a single




FIG. 5: Variation of allowed parameter space of the real coupling $c_L^{S',bq} = c_R^{S',bq}$ with scaling dimension d_U for B_d mixing (left) and B_s mixing (right). Black plotted points agree with the CP conserving mixing quantities $\Delta M_{d,s}$, while grey points also agree with the CP phases $\phi_{d,s}$.

real coupling, [13]. For this sub-case we can now set the bounds,

$$|c_L^{S',bd}| = |c_R^{S',bd}| \leq 0.00034 \quad (\Delta M_d \text{ only}) \quad (26)$$

$$|c_L^{S',bd}| = |c_R^{S',bd}| \leq 0.00012 \quad (\Delta M_d \& \phi_d) \quad (27)$$

$$|c_L^{S',bs}| = |c_R^{S',bs}| \leq 0.0013 \quad (\Delta M_s \text{ only}) \quad (28)$$

$$0.00058 \leq |c_L^{S',bs}| = |c_R^{S',bs}| \leq 0.0013 \quad (\Delta M_s \& \phi_s) \quad (29)$$

for $d_U = \frac{3}{2}$.




FIG. 6: Plot of the allowed $|c_L^{S',bq}| - \phi_L^{S',bq}$ parameter space for B_d mixing (left) and B_s mixing (right) for the case of a single complex coupling $c_L^{S',bq} \neq 0$ and $c_R^{S',bq} = 0$ and scaling dimension fixed as $d_U = 3/2$. Black plotted points agree with the CP conserving mixing quantities $\Delta M_{d,s}$, while grey points also agree with $\phi_{d,s}$.

The allowed parameter space for the third case, a single complex coupling, $c_L^{S',bq} = |c_L^{S',bq}| e^{i\phi_L^{S',bq}}$ and $c_R^{S',bq} = 0$, is

shown in fig. 6. In this case we have the bounds,

$$|c_L^{S',bd}| \leq 0.00029 \quad (\Delta M_d \text{ only}) \quad (30)$$

$$|c_L^{S',bd}| \leq 0.00018 \quad (\Delta M_d \& \phi_d) \quad (31)$$

$$|c_L^{S',bs}| \leq 0.0014 \quad (\Delta M_s \text{ only}) \quad (32)$$

$$0.00046 \leq |c_L^{S',bs}| \leq 0.0014 \quad \text{and} \quad 48.2^\circ \leq \phi_L^{S',bs} \leq 115.6^\circ \quad (\Delta M_s \& \phi_s) \quad (33)$$




FIG. 7: Plot of the allowed $|c_L^{S',bq}|$ - $|c_R^{S',bq}|$ parameter space for B_d mixing(left) and B_s mixing(right) for the case of two complex couplings $c_L^{S',bq} \neq 0$ and $c_R^{S',bq} \neq 0$. Black plotted points agree with the CP conserving quantities $\Delta M_{d,s}$, while grey points also agree with $\phi_{d,s}$.

For the final case of two complex couplings, $c_L^{S',bq} = |c_L^{S',bq}| e^{i\phi_L^{S',bq}}$ and $c_R^{S',bq} = |c_R^{S',bq}| e^{i\phi_R^{S',bq}}$, we again consider two sub-cases, $c_L^{S',bq} \neq c_L^{S',bq}$ and $c_L^{S',bq} = c_L^{S',bq}$. The first sub-case, $c_L^{S',bq} \neq c_L^{S',bq}$, is shown in fig. 7 where the allowed parameter space is again stretched along two lines. This is again similar to the parameter space found for the operators $\partial^\mu \mathcal{O}_U$ and $\mathcal{O}_U^\mu$. From the left panel of fig. 7 we can see that again the inclusion of the CP constraint from ϕ_d decreases the allowed region further. The right panel shows that CP phase ϕ_s has the same effect and further disfavors a small region near the origin. Fig. 8 displays the allowed parameter space for the sub-case $c_L^{S',bq} = c_L^{S',bq}$ where we can impose the bounds,

$$|c_L^{S',bd}| = |c_R^{S',bd}| \leq 0.00037 \quad (\Delta M_d \text{ only}) \quad (34)$$

$$|c_L^{S',bd}| = |c_R^{S',bd}| \leq 0.00022 \quad (\Delta M_d \& \phi_d) \quad (35)$$

$$|c_L^{S',bs}| = |c_R^{S',bs}| \leq 0.0017 \quad (\Delta M_s \text{ only}) \quad (36)$$

$$0.00056 \leq |c_L^{S',bs}| = |c_R^{S',bs}| \leq 0.0017 \quad \text{and} \quad -41.8^\circ \leq \phi_L^{S',bs} \leq 25.6^\circ \quad (\Delta M_s \& \phi_s) \quad (37)$$

for a scaling dimension of $d_U = 3/2$

IV. CONCLUSION

It was recently suggested in [27] that conformal invariance implies constraints on the scaling dimension, $1 \leq d_U \leq 2$, and $3 \leq d_U \leq 4$, for scalar and vector unparticles respectively. These constraints imply a rather large suppression of the contribution of vector and non-primary scalar unparticles to $B_{d,s}$ mixing, except for in a rather small range of scaling dimension close to integer values. The largest range where non-negligible contributions exist is for the non-primary operator $\partial^\mu \mathcal{O}_U$ with $2 \leq d_U \lesssim 2.1$. For primary scalar unparticles the suppression is far less and as such the range of scaling dimension for which sizable contributions to $B_{d,s}$ mixing exists is far larger. As a result, we have here considered, for the first time, the contribution of a primary scalar unparticle operator $\mathcal{O}_U$ to $B_{d,s}$ mixing. Considering a number of different coupling patterns, we have determined the allowed parameter space and set bounds on the unparticle couplings in each case. To illustrate the impact of the inclusion of the constraints from the CP




FIG. 8: Plot of the allowed $|c_L^{S',bq}| - \phi_L^{S',bq}$ parameter space for B_d mixing(left) and B_s mixing(right) for the case of one complex coupling $c_L^{S',bq} = c_R^{S',bq}$ and scaling dimension fixed as $d_U = 3/2$. Black plotted points agree with the CP conserving mixing quantities $\Delta M_{d,s}$, while grey points also agree with $\phi_{d,s}$.

phases associated with $B_{d,s}$ mixing, in particular the recently measured hint of new physics in ϕ_s , we have analyzed constraints with and without the inclusion of $\phi_{d,s}$. In each case we have found that these CP phases have an important role to play in constraining the parameter space, highlighted by the case of a single real unparticle coupling, $c_L^{S',bs} \neq 0$ and $c_R^{S',bs} = 0$, which was excluded for all values of the scaling dimension d_U .

-
- [1] H. Georgi, Phys. Lett. B **650** (2007) 275 [arXiv:0704.2457 [hep-ph]].
 - [2] H. Georgi, Phys. Rev. Lett. **98**, 221601 (2007) [arXiv:hep-ph/0703260].
 - [3] T. Banks and A. Zaks, Nucl. Phys. B **196** (1982) 189.
 - [4] K. Cheung, W. Y. Keung and T. C. Yuan, Phys. Rev. Lett. **99** (2007) 051803 [arXiv:0704.2588 [hep-ph]].
 - [5] M. Luo and G. Zhu, Phys. Lett. B **659** (2008) 341 [arXiv:0704.3532 [hep-ph]].
 - [6] C. H. Chen and C. Q. Geng, Phys. Rev. D **76** (2007) 115003 [arXiv:0705.0689 [hep-ph]].
 - [7] X. Q. Li and Z. T. Wei, Phys. Lett. B **651** (2007) 380 [arXiv:0705.1821 [hep-ph]].
 - [8] R. Mohanta and A. K. Giri, Phys. Rev. D **76** (2007) 075015 [arXiv:0707.1234 [hep-ph]].
 - [9] A. Lenz, Phys. Rev. D **76** (2007) 065006 [arXiv:0707.1535 [hep-ph]].
 - [10] S. L. Chen, X. G. He, X. Q. Li, H. C. Tsai and Z. T. Wei, arXiv:0710.3663 [hep-ph].
 - [11] R. Mohanta and A. K. Giri, Phys. Lett. B **660** (2008) 376 [arXiv:0711.3516 [hep-ph]].
 - [12] C. H. Chen, C. S. Kim and Y. W. Yoon, arXiv:0801.0895 [hep-ph].
 - [13] J. K. Parry, arXiv:0806.4350 [hep-ph].
 - [14] R. Zwicky, Phys. Rev. D **77** (2008) 036004 [arXiv:0707.0677 [hep-ph]].
 - [15] A. Abulencia *et al.* [CDF Collaboration], Phys. Rev. Lett. **97** (2006) 242003 [arXiv:hep-ex/0609040].
 - [16] V. M. Abazov *et al.* [D0 Collaboration], arXiv:0802.2255 [hep-ex].
 - [17] T. Aaltonen *et al.* [CDF Collaboration], arXiv:0712.2397 [hep-ex].
 - [18] M. Bona *et al.* [UTfit Collaboration], arXiv:0803.0659 [hep-ph].
 - [19] A. Lenz and U. Nierste, JHEP **0706** (2007) 072 [arXiv:hep-ph/0612167].
 - [20] M. Bona *et al.* [UTfit Collaboration], JHEP **0803** (2008) 049 [arXiv:0707.0636 [hep-ph]].
 - [21] D. Becirevic *et al.*, Nucl. Phys. B **634** (2002) 105 [arXiv:hep-ph/0112303].
 - [22] A. J. Buras, M. Jamin and P. H. Weisz, Nucl. Phys. B **347** (1990) 491.
 - [23] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963).
 - [24] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
 - [25] *et al.* [Tevatron Electroweak Working Group and CDF Collaboration and D0 Collab], arXiv:0808.1089 [hep-ex].
 - [26] T. Inami and C. S. Lim, Prog. Theor. Phys. **65** (1981) 297 [Erratum-ibid. **65** (1981) 1772].
 - [27] B. Grinstein, K. A. Intriligator and I. Z. Rothstein, Phys. Lett. B **662**, 367 (2008) [arXiv:0801.1140 [hep-ph]].
 - [28] G. Mack, Commun. Math. Phys. **55** (1977) 1.
 - [29] It should be noted that the combination of $D\bar{D}$ and CDF results for ϕ_s were made without full knowledge of the likelihoods, which could have an effect on the significance of this SM deviation. These are now being made available and a new experimental combination is due to be released soon. Until that time this present hint of new physics is the best available and remains a very exciting prospect.